

SUBGROUPS OF DEPTH THREE AND MORE

Sebastian Burciu¹ and Lars Kadison²

¹ *Inst. of Math. “Simion Stoilow” of the Romanian Academy P.O. Box 1-764,
Bucharest, Romania, RO-014700*

e-mail: sebastian.burciu@imar.ro

² *Department of Mathematics, University of Pennsylvania, David Rittenhouse Lab, 209 S. 33rd St.
Philadelphia, PA 19104*

e-mail: lkadison@math.upenn.edu

Abstract

A subalgebra pair of semisimple complex algebras $B \subseteq A$ with inclusion matrix M is depth two iff $MM^t M \leq nM$ for some positive integer n and all corresponding entries. If A and B are the group algebras of finite group-subgroup pair $H < G$, the induction-restriction table equals M and $S = MM^t$ satisfies $S^2 \leq nS$ iff the subgroup H is depth three in G ; similarly depth $n > 3$ by successive right multiplications of this inequality with alternately M and M^t . We show that a Frobenius complement in a Frobenius group is a nontrivial class of examples of depth three subgroups. A tower of Hopf algebras $A \supseteq B \supseteq C$ is shown to be depth-3 if $C \subset \text{core}(B)$; and this is also a necessary condition if A , B and C are group algebras.

INTRODUCTION

Induction of characters from a subgroup to a group is a useful technique for completing character tables [8] found by nineteenth century algebraists. At about the same time, Frobenius discovered reciprocity, which in modern terms states that induction is naturally isomorphic to coinduction of G -modules, either forming an adjoint pair with the restriction functor, and applies to any Frobenius extension of algebras.

Finite index subfactors are a certain type of Frobenius extension, where an analytic notion of finite depth was discovered in connection with classification, with depth two being part of a remarkable type of Galois theory of paragroups. The notion of finite depth was eventually made algebraic and applied to Frobenius extensions; later, depth two and its Galois theory of quantum groupoids and Hopf algebroids were exposed in simplest terms for ring extensions (see [18] for an application to J. Roberts field algebra construction [16]).

It was noted in [10] that the notion of depth two applies to characters of a finite group and subgroup pair via complex group algebras: a subgroup is depth two if no new constituents arise when inducing-restricting-inducing a character as compared with inducing just one time. By means of general theory in one direction and Mackey theory in the other, depth two subgroup is shown to be precisely a normal subgroup [10]. (A similar statement is true for semisimple Hopf $\mathbb{C}$ -subalgebras [4].) In this paper we generalize this approach to depth two subgroup to a semisimple subalgebra pair, giving a condition in terms of inclusion matrix [7], which is the same as a induction-restriction table [1] up to a permutation change of basis. The depth two

condition is essentially that the cube of the inclusion matrix is less entrywise than a multiple of the inclusion matrix, noted more precisely in the abstract and Proposition 1.2 below.

In [11] it was shown that finite depth Frobenius extension has a simplified definition in terms of a generalization of depth two to a tower of three algebras in the Jones tower. In this paper we extend a particular case of an embedding theorem in [11] to characterization of certain finite depth separable Frobenius extension in terms of depth two extension in Jones tower (see Theorems 2.1 and 2.5 below). Then one may check that a subgroup is depth three or more by comparing cube of symmetric matrix S of inner products of induced irreducible characters with multiples of S (see Prop. 2.2). In somewhat the same spirit, Corollary 2.9 below implies that a subgroup is depth three if no new constituents arise from applying restriction-induction one extra time to a character.

Although amusing to test for depth three property from character tables of groups and non-normal subgroup, it is not clear from this definition what precisely a depth three subgroup is. A number of proposals to remedy this are given below: depth three quasi-bases are given in Theorem 2.10, a characterization of certain depth three Frobenius extension in terms of similar bimodules, tensor-square and overalgebra in Theorem 2.7, and a class of examples in Section 3, a Frobenius group and its Frobenius complement. Even the notion of depth-3 tower of algebras may be viewed as an alternative to defining finite depth in terms of iterated endomorphism algebra extensions (perhaps applied instead to an iteration of another useful construction). Depth-3 towers of finite group algebras are completely classified in Theorem 1.1 following the spirit of [11]. Depth-3 towers of Hopf algebras are also considered at the end of the second section. A tower of Hopf algebras $A \supseteq B \supseteq C$ is depth-3 if $C \subseteq \text{core}(B)$ (see subsection 1.6 for the definition of the core of a Hopf subalgebra). Using then notion of kernel of a module introduced in [3] we formulate a conjecture on the *core* of a Hopf subalgebra. This conjecture would imply that the condition $C \subseteq \text{core}(B)$ is also a necessary condition for the Hopf algebra tower $A \supseteq B \supseteq C$ to be depth-3 (which is true for group algebras by the Theorem 1.1 below). Although our algebras are often over the complex numbers, the paper is hopefully written in a change-of-characteristic-friendly way.

1 PRELIMINARIES ON DEPTH TWO EXTENSIONS

All algebras in this paper are associative algebras (not necessarily commutative) over a field k . Given an (A, A) -bimodule M , we let M^A denote the A -central elements $\{m \in M \mid \forall a \in A, am = ma\}$.

Two $r \times s$ matrices M and N of non-negative integers satisfy $M \leq N$ if each of the coefficients $m_{ij} \leq n_{ij}$: this property is independent of permutation of bases. Note that if X is a third $q \times r$ matrix of non-negative integers, then $XM \leq XN$; if X is $s \times q$, then $MX \leq NX$. We say M is strictly positive if all entries $m_{ij} > 0$.

1.1 FROBENIUS EXTENSIONS

A Frobenius extension $A|B$ is an extension of associative algebras where the natural bimodule ${}_B A_A$ is isomorphic to the (B, A) -bimodule $\text{Hom}(A_B, B_B)$ (of right B -module homomorphisms) given by $(b \cdot f \cdot a)(x) = bf(ax)$ for $a, x \in A, b \in B, f \in \text{Hom}(A_B, B_B)$. This is equivalent to the existence of a mapping $F \in \text{Hom}({}_B A_B, {}_B B_B)$ with dual bases $\{x_i\}_{i=1}^n$ and $\{y_i\}_{i=1}^n$ such that $\sum_{i=1}^n F(ax_i)y_i = a$ and $\sum_{i=1}^n x_i F(y_i a) = a$ for all $a \in A$: we call the data system F a Frobenius homomorphism with dual bases $\{x_i\}, \{y_i\}$.

For example, a group algebra $A = k[G]$ is a Frobenius extension of any subgroup algebra $B = k[H]$, where $H \leq G$ is a subgroup of finite index $[G : H] = n$. For if $\{g_i\}_{i=1}^n$ denotes left coset representatives of H in G , where $g_1 = 1_G$, a Frobenius system is given by $x_i = g_i^{-1}, y_i = g_i$ with bimodule projection (then also split extension) given by $(n_{hg_i} \in k)$

$$F\left(\sum_{i=1}^n \sum_{h \in H} n_{hg_i} h g_i\right) = \sum_{h \in H} n_h h, \quad (1)$$

a routine exercise.

A Frobenius extension $A|B$ enjoys isomorphic tensor-square and endomorphism ring as (A, A) -bimodules. We note that $A \otimes_B A \cong \text{End}_B A$ via $x \otimes_B y \mapsto \lambda(x) \circ F \circ \lambda(y)$. Also $A \otimes_B A \cong \text{End}_B A$ via $x \otimes y \mapsto \rho(y) \circ F \circ \rho(x)$ [9]. Composing the two isomorphisms we obtain an anti-isomorphism $\text{End}_B A \rightarrow \text{End}_B A$ given by $f \mapsto \sum_i F(-f(x_i))y_i$, which restricts to an anti-automorphism on the subring $\text{End}_{BA} A$, and plays the role of antipode in case of depth two Frobenius extension defined below.

1.2 SEPARABLE EXTENSIONS

If the characteristic of the ground field k is coprime to $[G : H] = n$, then the extension of group algebras $A|B$ noted above is a separable extension: i.e., the multiplication map $\mu : A \otimes_B A \rightarrow A$ is a split (A, A) -epimorphism. The image of 1_A under a section $A \rightarrow A \otimes_B A$ is a separability element $e = \sum_{i=1}^n e_i \otimes_B f_i$ satisfying $ae = ea$ for all $a \in A$ and $\mu(e) = \sum_{i=1}^n e_i f_i = 1_A$, which characterizes separable extension. Notice that

$$\frac{1}{[G : H]} \sum_{i=1}^n g_i^{-1} \otimes_B g_i \quad (2)$$

is a separability element for the group algebras A over B .

In the situation that $C \supseteq A \supseteq B$ is a tower of algebras and $A|B$ is a separable extension, the canonical epi $C \otimes_B C \rightarrow C \otimes_A C$ given by $c_1 \otimes_B c_2 \mapsto c_1 \otimes_A c_2$ splits. A section for this mapping is of course given by $c_1 \otimes_A c_2 \mapsto \sum_{i=1}^n c_1 e_i \otimes_B f_i c_2$.

1.3 DEPTH-3 TOWERS OF ALGEBRAS

A tower of three algebras $A \supseteq B \supseteq C$, where C is a unital subalgebra of B which is in turn unital subalgebra of A , is said to be *right depth-3*, or *right d-3*, if there is a complementary (A, C) -bimodule P and $n \in \mathbb{N}$ such that

$$A \otimes_B A \oplus P \cong A^n \quad (3)$$

as natural (A, C) -bimodules. Equivalently, there is a split (A, C) -bimodule epimorphism from a finite direct sum of A with itself to $A \otimes_B A$ (P is the kernel of such an epi).

Left d-3 towers are defined oppositely, so that $A \supseteq B \supseteq C$ is left d-3 iff the tower of opposite algebras $A^{\text{op}} \supseteq B^{\text{op}} \supseteq C^{\text{op}}$ is right d-3. It has been noted in [11, 5] that if $A|B$ is a Frobenius, or quasi-Frobenius (QF, where isomorphisms above are replaced by similarity of bimodules) extension, then left d-3 is equivalent with right d-3 extension.

1.3.1 Depth-3 towers of semisimple algebras

Suppose a tower $A \supseteq B \supseteq C$ of semisimple finite dimensional k -algebras is right d-3. Tensoring eq. (3) by $-\otimes_C M$, we obtain the following inequality:

$$\langle M \uparrow_C^A \downarrow_B^A \uparrow_B^A, Q \rangle \leq n \langle M \uparrow_C^A, Q \rangle$$

which holds for any simple C -module M and any simple A -module Q .

Using this relation a necessary and sufficient condition for a tower of groups to be depth-3 will be given in the next theorem. For H a subgroup of G let

$$\text{core}_G(H) = \bigcap_{g \in G} {}^g H$$

be the largest subgroup of H which is normal in G . (Here ${}^g H = gHg^{-1}$.)

Let $G \supseteq N \supseteq H$ be a tower of groups. Since the normal closure H^G is the subgroup of G generated by the elements ghg^{-1} with $g \in G$ and $h \in H$ note that $H \subseteq \text{core}_G(N)$ if and only if $H^G \subseteq N$.

Theorem 1.1. *A tower $G \supseteq N \supseteq H$ of groups is depth-3 if and only if $H \subset \text{core}_G(N)$.*

Proof. If $H \subset \text{core}_G(N)$ then $H^G \subseteq N$ and the proof of Theorem 3.1 from [11] applies.

Suppose now that the tower is depth-3. The above argument for the tower $kG \supseteq kN \supseteq kH$ of semisimple algebras implies that there is $n \in \mathbb{N}$ such that

$$\langle \alpha \uparrow_H^G \downarrow_N^G \uparrow_N^G, \mu \rangle \leq n \langle \alpha \uparrow_H^G, \mu \rangle$$

for any characters $\alpha \in \text{Irr}(H)$ and $\mu \in \text{Irr}(G)$.

Put $\mu = 1_G$, the trivial character in the above inequality. Since $\langle \alpha \uparrow_H^G, 1_G \rangle = \langle \alpha, 1_H \rangle$ it follows that $\langle \alpha \uparrow_H^G \downarrow_N^G \uparrow_N^G, 1_G \rangle = 0$ if $\alpha \neq 1_H$. By Frobenius reciprocity this implies that $\langle \alpha \uparrow_H^G \downarrow_N^G, 1_N \rangle = 0$ if $\alpha \neq 1_H$.

On the other hand applying Mackey's theorem one has:

$$\begin{aligned} 0 = \langle \alpha \uparrow_H^G \downarrow_N^G, 1_N \rangle &= \sum_{NgH \in N \backslash G/H} \langle {}^g \alpha \downarrow_{N \cap {}^g H}^{{}^g H} \uparrow_{N \cap {}^g H}^N, 1_N \rangle \\ &= \sum_{NgH \in N \backslash G/H} \langle {}^g \alpha \downarrow_{N \cap {}^g H}^{{}^g H}, 1_{N \cap {}^g H} \rangle \\ &= \sum_{NgH \in N \backslash G/H} \langle \alpha \downarrow_{g^{-1}N \cap H}^H, 1_{g^{-1}N \cap H} \rangle \\ &= \sum_{NgH \in N \backslash G/H} \langle \alpha, 1_{g^{-1}N \cap H} \uparrow_{g^{-1}N \cap H}^H \rangle \end{aligned}$$

On the other hand using Frobenius reciprocity again one has

$$\langle 1_H, 1_{g^{-1}N \cap H} \uparrow_{g^{-1}N \cap H}^H \rangle = \langle 1_{g^{-1}N \cap H}, 1_{g^{-1}N \cap H} \rangle = 1$$

Thus

$$1_{g^{-1}N \cap H} \uparrow_{g^{-1}N \cap H}^H = 1_H$$

which implies that $H = g^{-1}N \cap H$ or $H \subset g^{-1}N = g^{-1}Ng$. Thus $H \subset \text{core}_G(N)$. $\square$

1.4 DEPTH TWO ALGEBRA EXTENSIONS

An algebra extension $A \supseteq B$ is defined to be *right depth two* (equivalently, subalgebra $B \subseteq A$ is rD2) if the partially trivial tower $A \supseteq B \supseteq B$ is right d-3; similarly we define left D2 in terms of partially trivial left d-3 tower.

It is obvious that a finite dimensional algebra A is a depth two extension of its unit subalgebra $B = k1_A$: if $\dim_k A = n$, then of course ${}_A A \otimes_k A \cong {}_A A^n$. Similarly, we may show that if C is a finite dimensional algebra, the tensor algebra $A = C \otimes B$ is a depth two extension of its subalgebra $B = 1_C \otimes B$.

The main examples in the literature of depth two extension are Hopf-Galois extensions as well as its classical, weakened and pseudo- variants.

The defining Condition (3), with $B = C$, for right depth two extension is similar to the characterization of projective module as isomorphic to a direct summand of a free module. Like the derivation of projective bases for a projective module, we may derive from this condition right D2 quasi-bases for the right D2 extension $A|B$ as follows. For any ring extension, using the hom-tensor relation, note that $\text{Hom}({}_A A \otimes_B A_B, {}_A A_B) \cong \text{End}_B A_B$. By evaluation at 1_A note that $\text{Hom}({}_A A_B, {}_A A \otimes_B A_B) \cong (A \otimes_B A)^B$.

Then the split epi from $\pi: A^n \rightarrow A \otimes_B A$ satisfies an equation $\pi \circ \sigma = \text{id}_{A \otimes_B A}$. We have n standard split epis $A^n \rightarrow A$, which compose with π and σ to give the equation $\sum_{i=1}^n f_i \circ g_i = \text{id}_{A \otimes_B A}$, where $f_i \in \text{Hom}(A, A \otimes_B A)$ and $g_i \in \text{Hom}(A \otimes_B A, A)$, to which we apply the simplifications noted above. Suppose $f_i \mapsto u_i \in (A \otimes_B A)^B$, while $g_i \mapsto \gamma_i \in \text{End}_B A_B$ for each $i = 1, \dots, n$. As a consequence, we obtain for any $x, y \in A$ the identity

$$x \otimes_B y = \sum_{i=1}^n x \gamma_i(y) u_i \quad (4)$$

Note that an extension $A|B$ having elements $u_i \in (A \otimes_B A)^B$ and endomorphisms $\gamma_i \in \text{End}_B A_B$ satisfying this identity, eq. (4), also implies that $A|B$ is right D2, since $A^n \rightarrow A \otimes_B A$ given by $(a_1, \dots, a_n) \mapsto \sum_i a_i u_i$ is an (A, B) -epimorphism with section given by $x \otimes_B y \mapsto (x \gamma_1(y), \dots, x \gamma_n(y))$. For example, a normal subgroup N of index n in any group G (over any ground ring) is depth two with D2 quasi-bases given by $u_i = g_i^{-1} \otimes g_i$ and $\gamma_i(g) = F(g g_i^{-1}) g_i$ for coset representatives $\{g_1 = e, g_2, \dots, g_n\}$.

1.5 WHEN INCLUSION MATRIX IS DEPTH TWO

Let the ground field $k = \mathbb{C}$ be the complex numbers when we consider semisimple algebras, which consequently become multi-matrix algebras (or split semisimple algebras). Suppose $B \subseteq A$ is a subalgebra pair of semisimple algebras. As one constructs an induction-restriction table for a subgroup H in a finite group G [1, p. 166], we briefly review the procedure for generalizing to any pair of semisimple algebras (such as finite dimensional complex group algebras). Label the simples of A by $V_1, \dots, V_s$ and the simple modules of B by $W_1, \dots, W_r$. To obtain the i 'th column restrict the i 'th simple A -module V_i to a B -module and express in terms of direct sum of simples

$$V_j \downarrow_B \cong \oplus_{i=1}^r m_{ij} W_i \quad (5)$$

We let M be the $r \times s$ -matrix, or table, with entries m_{ij} : $M = (m_{ij})$. By a well-known generalization of Frobenius reciprocity, the rows give induction of the B -simples:

$$W_i \uparrow^A = W_i^A = \oplus_{j=1}^s m_{ij} V_j \quad (6)$$

since $W_j^A = W_j \otimes_B A$ and $V_i \downarrow_B \cong \text{Hom}(A_B, V_i)$; i.e., if $[W_j^A, V_i]$ denotes the number of constituents in W_j^A isomorphic to V_i , Frobenius reciprocity is given by

$$[W_i^A, V_j] = m_{ij} = [W_i, V_j \downarrow_B] \quad (7)$$

The matrix M is also known as the inclusion matrix of B in A [7].

For example, the induction-restriction table (based on Frobenius reciprocity $(\psi_i^G, \chi_j)_G = (\psi_i, \chi_j \downarrow_H)_H$) for the standard embedding of permutation groups $S_2 \leq S_3$ is given by

$$\begin{array}{c|ccc} S_2 \leq S_3 & \chi_1 & \chi_2 & \chi_3 \\ \hline \psi_1 & 1 & 0 & 1 \\ \psi_2 & 0 & 1 & 1 \end{array} \quad M = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \end{pmatrix} \quad \begin{array}{ccc} \overset{1}{\bullet} & \overset{2}{\bullet} & \overset{1}{\bullet} \\ & \backslash & / \\ & \overset{\circ}{1} & \overset{\circ}{1} \end{array}$$

where $\psi_1 = 1_H$, $\chi_1 = 1_G$ denote the trivial characters, ψ_2, χ_2 the sign characters, and χ_3 the two-dimensional irreducible character of S_3 . Note too the inclusion diagram or Bratteli diagram, a bicolored weighted multigraph [7].

For example, $1_H^G = \chi_1 + \chi_3$ and $1_H^G \downarrow_H = 2 \cdot 1_H + \psi_2$. Burciu [3] notes that a subgroup H is normal in G if and only if $1_H^G \downarrow_H = [G : H]1_H$. In [10] it is established that the notion of depth two subalgebra for subalgebra pair of complex group algebras is equivalent to the notion of normal subgroup.

Proposition 1.2. *The inclusion matrix M of a subalgebra pair of semisimple complex algebras $B \subseteq A$ satisfies*

$$MM^t M \leq nM \quad (8)$$

for some positive integer n if and only if B is depth two subalgebra of A .

Proof. ($\Leftarrow$) The depth two condition $A \otimes_B A \oplus P \cong A^n$ as natural B - A -bimodules, becomes

$$[W_i^A \downarrow_B \uparrow^A, V_j] \leq n[W_i^A, V_j] = nm_{ij} \quad (9)$$

for all $i = 1, \dots, r$ and $j = 1, \dots, s$. But W_i^A is given by row i of M , or $e_i M$, where e_i denotes row matrix with all zeroes except 1 in i 'th column. Then $W_i^A \downarrow_B$ is given by $M(e_i M)^t = MM^t e_i^t$. Finally $W_i^A \downarrow_B \uparrow^A$ is given by $(MM^t e_i^t)^t M$, i.e. row i of $MM^t M$.

($\Rightarrow$) If the inclusion matrix M of semisimple subalgebra pair $B \subseteq A$ satisfies $MM^t M \leq nM$ for some $n \in \mathbb{Z}_+$, then $[\text{Ind}_B^A \text{Res}_B^A \text{Ind}_B^A W_i, V_j] \leq n[\text{Ind}_B^A W_i, V_j]$ for all B -simples W_i and A -simples V_j (fix these orderings). Via unique module decomposition into simples, we find a monic natural transformation $\text{Ind}_B^A \text{Res}_B^A \text{Ind}_B^A \hookrightarrow n\text{Ind}_B^A$ from category $B\text{-Mod}$ into $A\text{-Mod}$. Now B, A and so $B^{\text{op}} \otimes A$ are separable $\mathbb{C}$ -algebras, so as in [10, Theorem 2.1(6), pp. 3107-3108], we apply the natural monic to the right regular module B_B , apply the natural transformation property to all left multiplications λ_b ($b \in B$), and note that $A \otimes_B A \hookrightarrow A^n$ splits by Maschke as B - A -bimodule monic. Hence A is depth two over its subalgebra B . $\square$

1.6 DEPTH-3 TOWER OF OF HOPF ALGEBRAS

For $B \subset A$ an extension of finite dimensional Hopf algebras, define $\text{core}(B)$ to be the largest Hopf subalgebra of B which is normal in A . It is easy to see that $\text{core}(B)$ always exists (see also [3]). If $H \subset G$ is a group inclusion with $A = kG$ and $B = kH$ note that $\text{core}(B) = k\text{core}_G(H)$.

Theorem 1.3. Suppose that $A \supseteq B \supseteq C$ is a tower of semisimple Hopf algebras. If $C \subset \text{core}(B)$ then the tower is depth-3.

Proof. Since $\text{core}(B)$ is a normal Hopf subalgebra of A it follows that the extension $\text{core}(B) \subset A$ is D2 and therefore $A \otimes_{\text{core}(B)} A$ is a direct summand of the bimodule ${}_A(A^n)_{\text{core}(B)}$. Thus $A \otimes_{\text{core}(B)} A$ is also a direct summand of the $A - C$ bimodule ${}_A(A^n)_C$ since $C \subset \text{core}(B)$. Since $\text{core}(B) \subset B$ the canonical map

$$A \otimes_{\text{core}(B)} A \rightarrow A \otimes_B A$$

is a surjective morphism of $A - A$ -bimodules, in particular of $A - C$ bimodules. Since the category of $A \otimes C^{op}$ -modules is semisimple it follows that $A \otimes_B A$ is a direct summand in ${}_A(A^n)_C$. $\square$

1.6.1 Kernel of a module

Let A be a semisimple Hopf algebra over an algebraically closed field k . Then A is also cosemisimple and $S^2 = \text{Id}$ (see [13]). Let A_χ be the idempotent integral of A . Denote by $\text{Irr}(A)$ the set of irreducible A -characters and let $C(A)$ be the character ring of A with basis $\text{Irr}(A)$. There is an involution “ $*$ ” on $C(A)$ determined by the antipode.

Remark 1.4. If $X \subset C(A^*)$ is closed under multiplication then it generates a subbialgebra of A denoted by A_X [15]. Moreover if X is also closed under “ $*$ ” it follows from the same paper that A_X is a Hopf subalgebra. Since A is finite dimensional any subbialgebra is a Hopf subalgebra and therefore any subset X closed under multiplication is also closed under “ $*$ ”.

Let M be an A -module with character χ . Define $\ker_M$ to be the set of simple subcoalgebras C of A such that $cm = \varepsilon(c)m$ for all $c \in C$. It can be proven that the set $\ker_M$ is closed under multiplication and “ $*$ ” and therefore from [15] it generates a Hopf subalgebra A_M (or A_χ) of A [3]. One has $A_M = \bigoplus_{C \in \ker_M} C$.

Remark 1.5. 1) A_χ is the largest subbialgebra B of A such that $\chi \downarrow_B^A = \chi(1)\varepsilon_B$. Equivalently, A_χ is the largest subbialgebra B of A such that $AB^+A \subset \text{Ann}_A(M)$.

2) If $A = kG$ is a group algebra then $A_\chi = k[\ker \chi]$ where $\ker \chi$ is the kernel of the character χ .

3) It is not known if A_χ is a normal Hopf subalgebra of A . In [3] it was proven that A_χ is normal in A if $\chi \in Z(A^*)$.

4) If N is a submodule or a quotient of M then clearly $A_M \subset A_N$ (since A is semisimple).

Notation: If B is a Hopf subalgebra of A then we denote by $\varepsilon \uparrow_B^A$ the character $\varepsilon_B \uparrow_B^A$.

Proposition 1.6. Suppose B and C are Hopf subalgebras of a finite dimensional semisimple Hopf algebra A . If

$${}_A(A \otimes_B A)_C \oplus * \cong_A A^n{}_C \quad (10)$$

as A - C -bimodules, then

$$C \subset A_{\varepsilon \uparrow_B^A}.$$

Proof. As in subsection 1.3.1 it follows that

$$\langle M \uparrow_C^A \downarrow_B^A \uparrow_B^A, P \rangle \leq n \langle M \uparrow_C^A, P \rangle$$

for any simple left C -module M and simple A -module P .

In terms of the characters this can be written as

$$m_A(\alpha \uparrow_C^A \downarrow_B^A \uparrow_B^A, \chi) \leq n m_A(\alpha \uparrow_C^A, \chi) \quad (11)$$

for any irreducible character α of C and any irreducible character χ of A . Here m_A is the usual multiplication form on the character ring $C(A)$. Put $\chi = \varepsilon_A$, the trivial A -character, in the above inequality. Since $m_A(\alpha \uparrow_C^A, \varepsilon_A) = m_C(\alpha, \varepsilon_C)$ it follows that $m_A(\alpha \uparrow_C^A \downarrow_B^A \uparrow_B^A, \varepsilon_A) = 0$ if $\alpha \neq \varepsilon_C$. By Frobenius reciprocity this implies that $m_B(\alpha \uparrow_C^A \downarrow_B^A, \varepsilon_B) = 0$ if $\alpha \neq \varepsilon_C$. Adding over all irreducible characters $\alpha \in \text{Irr}(C)$ it follows that

$$m_B((\sum_{\alpha \in \text{Irr}(C)} \alpha(1)\alpha) \uparrow_C^A \downarrow_B^A, \varepsilon_B) = m_B(\varepsilon \uparrow_C^A \downarrow_B^A, \varepsilon_B) \quad (12)$$

Since $\sum_{\alpha \in \text{Irr}(C)} \alpha(1)\alpha$ is the regular character of C (see [14]) it follows that $(\sum_{\alpha \in \text{Irr}(C)} \alpha(1)\alpha) \uparrow_C^A \downarrow_B^A$ is the regular character of B multiplied by $\frac{|A|}{|C|}$. Thus $m_B(\varepsilon \uparrow_C^A \downarrow_B^A, \varepsilon_B) = \frac{|A|}{|B|}$. Frobenius reciprocity implies that $m_C(\varepsilon \uparrow_B^A \downarrow_C^A, \varepsilon_C) = \frac{|A|}{|B|}$. A dimension argument now shows that $\varepsilon \uparrow_B^A \downarrow_C^A = \frac{|A|}{|B|} \varepsilon_C$ and first item of Remark 1.5 implies that $C \subset A_{\varepsilon \uparrow_B^A}$. $\square$

The above Proposition and Theorem 1.3 suggest the following conjecture:

Conjecture 1. For any Hopf subalgebra B of a semisimple Hopf algebra A one has:

$$\text{core}(B) = A_{\varepsilon \uparrow_B^A}. \quad (13)$$

The next Proposition gives a description of $\text{core}(B)$ in terms of kernels and shows the inclusion $\text{core}(B) \subseteq A_{\varepsilon \uparrow_B^A}$. In order to prove it we need the following lemmas.

Lemma 1.7. *Let K and L be two Hopf subalgebras of a semisimple Hopf algebra A . If $\Lambda_K \Lambda_L = \Lambda_L$ then $K \subset L$.*

Proof. By Corollary 2.5 of [2] there is a coset decomposition for A

$$A = \oplus_{C/\sim} CL. \quad (14)$$

where $\sim$ is an equivalence relation on the set of simple subcoalgebras of A given by $C \sim C'$ if and only if $CL = C'L$. In [2] this equivalence relation is denoted by $r_{k,L}^A$. The equality $\Lambda_K \Lambda_L = \Lambda_L$ shows that any subcoalgebra of K is equivalent to $k1$ and therefore it is contained in L . $\square$

Lemma 1.8. *Suppose that B is a Hopf subalgebra of a semisimple finite dimensional Hopf algebra. Then $A_{\varepsilon \uparrow_B^A} \subset B$. Equality holds if and only if B is normal in A .*

Proof. Let $K = A_{\varepsilon \uparrow_B^A}$. By the definition of K it follows that AK^+ annihilates $A \otimes_B k$. On the other hand $A \otimes_B k \cong A/AB^+$ as A -modules and therefore $AK^+ \subset AB^+$. Thus $1 - \Lambda_K \in AB^+$ which implies $\Lambda_K \Lambda_B = \Lambda_B$. The above Lemma implies that $K \subset B$. The second statement of the lemma is Corollary 2.5 from [3]. $\square$

Proposition 1.9. *Suppose that B is a Hopf subalgebra of a semisimple finite dimensional Hopf algebra A . Define inductively*

$$B_0 = B, \quad B_{s+1} = A_{\varepsilon \uparrow_{B_s}^A}.$$

Then

$$B_1 \supseteq B_2 \supseteq \cdots \supseteq B_s \supseteq B_{s+1} \supseteq \cdots.$$

If $B_s = B_{s+1}$ then $B_s = \text{core}(B)$.

Proof. The above proposition implies that $B_r \supseteq B_{r+1}$ for any r . Since B is finite dimensional there is s such that $B_s = B_{s+1} = B_{s+2} = \dots$. Thus $B_s = A_{\varepsilon_{B_s}^A}$ and the above lemma implies that B is normal in A . We have to show that $\text{core}(B) = B_s$. Suppose that K is normal in A and that $K \subseteq B$. It is enough to show $K \subseteq B_s$. Clearly $K \subseteq B_0$. If $K \subseteq B_i$ then there is a canonical surjection of A -modules $A/AK^+ \rightarrow A/AB_i^+$. Thus $A_{\varepsilon_K^A} \subseteq A_{\varepsilon_{B_i}^A}$ by the last item of Remark 1.5. On the other hand $A_{\varepsilon_K^A} = K$ since K is normal. Therefore $K \subseteq B_{i+1}$. $\square$

1.6.2 The Correspondent of conjugate Hopf subalgebras

Let A be a semisimple Hopf algebra over an algebraically closed field k and let $\widehat{A^*}$ be the set of simple subcoalgebras of A . Since A is cosemisimple note that $\widehat{A^*}$ can be identified with $\text{Irr}(A^*)$ [12]. Let B be a Hopf subalgebra of A and C a simple subcoalgebra of A . Define

$$X_{C_B} = \{D \in \widehat{A^*} \mid dc\Lambda_B = \varepsilon(d)c\Lambda_B \text{ for all } c \in C, d \in D\}$$

Proposition 1.10. *The set X_{C_B} is closed under multiplication and “ $\cdot$ ” and it generates a Hopf subalgebra ${}^C B$ of A .*

Proof. By Remark 1.4 it is enough to show that the above set is closed under multiplication. Suppose that D and D' are subcoalgebras in X_{C_B} . If E is a simple subcoalgebra of DD' then any $e \in E$ can be written as $\sum_{i=1}^s d_i d'_i$ with $d_i \in D$ and $d'_i \in D'$. Then $ec\Lambda_B = \varepsilon(e)c\Lambda_B$ which show that $E \in X_{C_B}$. $\square$

Notation: ${}^C B$ will also be denoted with ${}^c B$ if c is the irreducible character of A^* corresponding to C .

Example 1.11. Let $A = kG$ and $B = kN$ where N is a subgroup of G . The simple subcoalgebras of A are kg with $g \in G$ and $\Lambda_B = \frac{1}{|N|} \sum_{n \in N} n$. Then ${}^g B = gBg^{-1}$ for all $g \in G$. Indeed $X_{gB} = \{h \in G \mid hg\Lambda_B = g\Lambda_B\} = \{h \in G \mid hgN = gN\} = gNg^{-1}$.

Proposition 1.12. *Let B be a Hopf subalgebra of A and $g \in G(A)$ be a grouplike element of A . Then ${}^g B = gBg^{-1}$.*

Proof. First note that ${}^1 B = B$. Clearly $B \subset {}^1 B$. On the other hand the definition of ${}^1 B$ implies that $\Lambda_{{}^1 B} \Lambda_B = \Lambda_B$. Then Lemma 1.7 implies ${}^1 B \subset B$.

Let now C be a simple subcoalgebra of ${}^g B$. Then $cg\Lambda_B = \varepsilon(c)g\Lambda_B$ for all $c \in C$. Thus $g^{-1}cg\Lambda_B = \varepsilon(c)\Lambda_B$ which shows that $g^{-1}Cg \subset {}^1 B = B$. Therefore $C \subset gBg^{-1}$ which shows that ${}^g B \subset gBg^{-1}$. A direct computations shows that $gBg^{-1} \subset {}^g B$. Thus ${}^g B = gBg^{-1}$. $\square$

Proposition 1.13. *Let B be a Hopf subalgebra of A . Then*

$$A_{\varepsilon_B^A} = \bigcap_{C \in \widehat{A^*}} {}^C B.$$

Proof. Recall the coset decomposition

$$A = \bigoplus_{C/\sim} CB. \tag{15}$$

form Corollary 2.5 of [2]. If k is the trivial B -module then

$$k \uparrow_B^A = \oplus_{C/\sim} {}^C B \otimes_B k.$$

From the definition of ${}^C B$ it follows that ${}^C B \otimes_B k$ is trivial as left ${}^C B$ -module. Therefore $\cap_{C \in \widehat{A}^*} {}^C B \subset A_{\varepsilon \uparrow_B^A}$.

Note that $k \uparrow_B^A = A \otimes_B k \cong A \Lambda_B$ as left A -modules via $a \Lambda_B \mapsto a \otimes_B \Lambda_B$. The decomposition 15 implies that ${}^C B \otimes_B k \cong C \Lambda_B$ under the above isomorphism. Any simple subcoalgebra of $A_{\varepsilon \uparrow_B^A}$ acts trivially on $k \uparrow_B^A$ and therefore on each ${}^C B \otimes_B k$. This implies that any such coalgebra is contained in ${}^C B$. Thus $A_{\varepsilon \uparrow_B^A} \subset \cap_{C \in \widehat{A}^*} {}^C B$ for any simple coalgebra $C \in \widehat{A}^*$. $\square$

Corollary 1.14. *Let B be a Hopf subalgebra of A . Then B is a normal Hopf subalgebra if and only if*

$$B = \cap_{C \in \widehat{A}^*} {}^C B.$$

Proof. Since $A_{\varepsilon \uparrow_B^A} = \cap_{C \in \widehat{A}^*} {}^C B$, this is Corollary 2.5 of [3]. $\square$

Remark 1.15. 1) Theorem 1.9 implies that $\text{core}(B) \subset A_{\varepsilon \uparrow_B^A} = \cap_{C \in \widehat{A}^*} {}^C B$. This can also be seen directly as follows. Fix $C \in \widehat{A}^*$. For any $x \in \text{core}(B)$ and $c \in C$ one has that $xc \Lambda_B = c_1(S(c_2)xc_3) \Lambda_B = c_1 \varepsilon(x) \varepsilon(c_2) \Lambda_B = \varepsilon(x) c \Lambda_B$ since $\text{core}(B)$ is normal in A . Thus $\text{core}(B) \subset {}^C B$.

2) If A_χ is normal Hopf algebra for any $\chi \in \text{Irr}(A)$ then Proposition 1.9 implies the above conjecture on the core of a Hopf subalgebra.

2 DEPTH THREE FROBENIUS EXTENSION

A Frobenius extension $A|B$ is defined to be *depth three* if the following tower of subalgebras in the endomorphism ring $E = \text{End} A_B$ is right or left depth-3: via the algebra monomorphism, left multiplication $\lambda : A \hookrightarrow E$ given by $\lambda(a)(x) = ax$ ($x, a \in A$) we obtain the (ascending) tower, $\lambda(B) \subseteq \lambda(A) \subseteq E$. By [11, Theorem 3.1] the given tower is left d-3 if and only if the tower is right d-3.

The definitions and first properties of depth two and three extensions are introduced in detail in [11]. There it is determined that a tower of three group algebras corresponding to the subgroup chain $G \geq H \geq K$ is depth-3 if the normal closure K^G (of K in G) is contained in H (and shown above in Theorem 1.1 to be a characterization of depth-3 tower of finite groups). In [10] it is shown that, with $k = \mathbb{C}$ and G a finite group, the group algebra A of G is depth two over subgroup algebra B of H if and only if H is a normal subgroup of G . This normality result for depth two subalgebras is extended to semisimple Hopf algebras over an algebraically closed field of characteristic zero in [4].

The following is a characterization of depth three for a separable, Frobenius extension in terms of the more familiar depth two property. The following is true more generally for QF-extensions [5, Theorem 3.8].

Theorem 2.1. *Suppose $A|B$ is a separable extension and Frobenius extension. Let E denote $\text{End} A_B$ and $\lambda : A \hookrightarrow E$ be understood as the extension $E|A$. The $A|B$ is depth three if and only if the composite extension $E|B$ is depth two.*

Sketch of Proof. ($\Rightarrow$) This direction does not apply separability. By the Frobenius extension property, we noted above that $E \cong A \otimes_B A$ as (A, A) -bimodules. Then $E \otimes_A E \otimes_A E \cong E \otimes_B E$ as natural (E, E) -bimodules. By definition of right D3 extension, $E \otimes_A E$ is isomorphic to direct summand of E^n as natural (E, B) -bimodules for some $n \in \mathbb{N}$, whence $E \otimes_A E \otimes_A E \cong E \otimes_B E$ is (E, B) -isomorphic to a direct summand of $E \otimes_A E^n$, which in turn is isomorphic to a direct summand of E^{n^2} by the right D3 property. Hence $E | B$ is right D2, since $E \otimes_B E \oplus * \cong E^{n^2}$ as natural (E, B) -bimodules.

($\Leftarrow$) There is a split (E, B) -epimorphism from $E^n \rightarrow E \otimes_B E$ for some $n \in \mathbb{N}$. In addition, there is a split (E, E) -epimorphism from $E \otimes_B E \rightarrow E \otimes_A E$ by the separability property of the extension $A | B$. Composing the two split epis we obtain a split epi $E^n \rightarrow E \otimes_A E$ showing $A | B$ is right D3. $\square$

The proposition below has a proof useful to the exposition, although the result is improved somewhat in subsection 2.1.

Proposition 2.2. *Let M be the inclusion matrix of a subalgebra pair of semisimple complex algebras $B \subseteq A$, and $S = MM^t$. The symmetric matrix S satisfies*

$$S^3 \leq nS \quad (16)$$

for some positive integer n if and only if B is a depth three subalgebra of A .

Proof. Let $M_m(\mathbb{C}) = \text{End}_{\mathbb{C}} A = \mathcal{E}$ where $m = \dim A$, which contains both A and B via left regular representation. It is shown in [7, 2.3.5] that the centralizers $\mathcal{E}^A \subseteq \mathcal{E}^B$ have transpose inclusion matrix; i.e. inclusion matrix of $A \hookrightarrow \text{End} A_B$ via $a \mapsto \lambda_a$ is M^t . It is not hard to show from transitivity of induction that matrix multiplication yields new inclusion matrix of two successive subalgebra pairs. Hence, inclusion matrix of $B \hookrightarrow E$ via $b \mapsto \lambda_b$ ($b \in B$) is given by MM^t .

The algebra A is separable, whence separable extension over B . The extension $A \supseteq B$ is a split Frobenius extension by application of [7, Goodman-De la Harpe-Jones, ch. 2], very faithful conditional expectations. Then A_B is a progenerator since B is semisimple and $B \hookrightarrow A$ is split B -module monic, so E and B are Morita equivalent semisimple algebras. By the theorem above, $B \subseteq A$ is depth three iff $B \hookrightarrow E$ is depth two, and we may apply Proposition 1.2 to the composite inclusion matrix $S = MM^t$. $\square$

In general for any subgroup H in finite group G with inclusion matrix M , if the irreducible characters of H are given by $\{\psi_1, \dots, \psi_r\} = \text{Irr}(H)$, note that the matrix $S = MM^t$ is given by

$$S = \begin{pmatrix} \langle \psi_1^G | \psi_1^G \rangle & \dots & \langle \psi_1^G | \psi_r^G \rangle \\ \dots & \dots & \dots \\ \langle \psi_r^G | \psi_1^G \rangle & \dots & \langle \psi_r^G | \psi_r^G \rangle \end{pmatrix}. \quad (17)$$

For example, we revisit the inclusion $S_2 < S_3$ analyzed above. Note that

$$S = MM^t = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}. \quad (18)$$

Since S is strictly positive (i.e. has only positive whole number entries), it is clear that there is positive integer n such that $S^3 \leq nS$.

The notation in the proposition above with finite dimensional complex group algebras $B = \mathbb{C}[H]$ and $A = \mathbb{C}[G]$ is continued in the next corollary:

Corollary 2.3. *The subgroup H is depth three in G if the matrix S is strictly positive.*

Another example: the standard inclusion of full permutation group algebras $B = \mathbb{C}[S_3] \hookrightarrow \mathbb{C}[S_4] = A$ has inclusion matrix (computed from character tables in e.g. [6]) and symmetric matrix:

$$M = \begin{pmatrix} 1 & 0 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 & 1 \\ 0 & 0 & 1 & 1 & 1 \end{pmatrix} \quad S = \begin{pmatrix} 2 & 0 & 1 \\ 0 & 2 & 1 \\ 1 & 1 & 3 \end{pmatrix} \quad S^3 = \begin{pmatrix} 15 & 7 & 21 \\ 7 & 15 & 21 \\ 21 & 21 & 43 \end{pmatrix}$$

It is clear that there is no positive integer n for which $S^3 \leq nS$, since S has zero entries but S^3 is strictly positive. We conclude that S_3 is not a depth three subgroup of S_4 . (Using the next theorem one computes that S_3 is a depth five subgroup of S_4 .)

2.1 HIGHER DEPTH

Recall from [11] that depth $n > 2$ is defined as follows. Begin with a Frobenius extension (or QF extension [5]) $B = A_{-1} \subseteq A = A_0$. Let $A_1 = \text{End} A_B$ and inductively $A_n = \text{End}(A_{n-1})_{A_{n-2}}$. By the Frobenius hypothesis and its endomorphism ring theorem, $A_n \cong A \otimes_B \cdots \otimes_B A$ ($n+1$ times A). Embedding $A_n \hookrightarrow A_{n+1}$ via left regular representation λ , we obtain a Jones tower of algebras,

$$B \hookrightarrow A \hookrightarrow A_1 \hookrightarrow \cdots \hookrightarrow A_n \hookrightarrow A_{n+1} \hookrightarrow \cdots$$

The subalgebra B in A is depth n if $A_{n-2} \supseteq A_{n-3} \supseteq B$ is a depth-3 tower defined above; infinite depth if there is no such positive integer n . Of course, this agrees with the definition of depth three subalgebra above. If B and A are semisimple complex algebras, $A \supseteq B$ becomes a split, separable Frobenius extension via the construction of a very faithful conditional expectation [7]. This type of extension has an endomorphism ring theorem [9], and enjoys transitivity, so that all extensions in this Jones tower are split, separable Frobenius extensions, and all algebras are semisimple by Morita's theorem (or Serre's theorem on global dimension). Indeed, all the odd A_n 's are Morita equivalent to B , while all the even A_n 's are Morita equivalent to A . The proof of the lemma below is similar to that of Prop. 1.2 and therefore omitted. (One notes that $\text{Ind}_C^A \cong \text{Ind}_B^A \text{Ind}_C^B$ is given by the rows of matrix NM .)

Lemma 2.4. *Suppose $C \subseteq B \subseteq A$ is a tower of semisimple algebras with inclusion matrices N and M respectively. Then the tower is depth-3 if and only if there is a positive integer n such that*

$$NMM^t M \leq nNM \tag{19}$$

Notice that Prop. 1.2 follows from letting $B = C$ and N equal the identity matrix of rank $\dim Z(B)$. Conversely, if $A \supseteq B$ is depth two, and C any subalgebra of B , then the lemma follows in this special case from Prop. 1.2 by multiplying the inequality there from the left by the inclusion matrix N of $C \subseteq B$.

Let n, m and q denote positive integers below.

Theorem 2.5. *Suppose $B \subseteq A$ is a subalgebra pair of semisimple algebras. Let M be the inclusion matrix and $S = MM^t$. If $n = 2m + 1$ then $A \supseteq B$ is depth n if and only if $S^{m+1} \leq qS^m$ for some q . If $n = 2m$, then $A \supseteq B$ is depth n if and only if $S^m M \leq qS^{m-1}M$ for some q .*

Proof. The proof follows from noting that if M is the inclusion matrix of $B \subseteq A$, then M^t is the inclusion matrix of $A \hookrightarrow A_1$, and S is the inclusion matrix of their composite $B \hookrightarrow A_1$. The proof now follows from applying the last lemma to the depth-3 tower $B \hookrightarrow A_{n-3} \hookrightarrow A_{n-2}$ in the even and odd case. $\square$

It is worth emphasizing that a depth n algebra extension is also depth $n + 1$ (so one might denote this as depth $\geq n$); in the special case of the theorem, this is seen by multiplying the given inequality from the right by the inclusion matrix M or M^t . Of course one should strive to use the least depth to one's knowledge. Let m be a positive integer and G a finite group in the next result on subgroups of finite depth.

Corollary 2.6. *Suppose $H < G$ is a subgroup with symmetric matrix S . If S^m is a strictly positive matrix, then H is a subgroup of depth $2m + 1$ in G .*

Proof. Applying the theorem we see $S^{m+1} \leq qS^m$ for some positive integer q since S^m is a strictly positive matrix. $\square$

For example, while $S_3 < S_4$ is not D3 subgroup, we note that S^2 is already strictly positive order 3 matrix, whence it is a depth five subgroup (and it may be checked that it is not depth four).

As another example of a more cautionary note, the symmetries of a square D_4 in S_4 has zero entries in all powers of its order 5 matrix $S = MM^t$. However, one computes that $S^2M \leq 4SM$, so that D_4 is a depth four subgroup of S_4 according to Theorem 2.5.

In a forthcoming paper it will be shown that after a permutation of the indices, the matrix S can be written as a sum of diagonal blocks. Moreover there is $p > 0$ such that the p -power of each diagonal block is a positive matrix. Applying Theorem 2.5 this implies that the extension $B \subseteq A$ is of depth $\geq 2p + 1$; in other words, all semisimple subalgebra pairs are of finite depth.

2.2 SIMPLIFIED CONDITION FOR DEPTH THREE

Again let $A \supseteq B$ be an algebra extension. In case A_B is a generator, such as when the extension is free or right split, there is a particularly simplified condition for when a Frobenius extension is depth three.

Theorem 2.7. *Suppose $A \supseteq B$ is a Frobenius extension where the natural module A_B is a generator. Then $A \supseteq B$ is depth three if and only if there is a B - B -bimodule P and positive integer n such that*

$${}_B A \otimes_B A_B \oplus P \cong {}_B A_B^n \quad (20)$$

Proof. ($\Rightarrow$) Let $E = \text{End} A_B$. By the Frobenius extension hypothesis on $A \supseteq B$, as E - A -bimodules $E \cong A \otimes_B A$ via the mapping in subsection 1.1. Recall that $A \supseteq B$ is depth three if $B \subseteq A \hookrightarrow E$ is depth three tower, i.e. ${}_E E \otimes_A E_B \oplus Q \cong {}_E E_B^n$ for some E - B -bimodule Q and positive integer n . Then by substitution

$${}_E A \otimes_B A \otimes_B A_B \oplus Q \cong {}_E A \otimes_B A_B^n. \quad (21)$$

But A_B is a progenerator by hypothesis, whence B and E are Morita equivalent algebras. The context bimodule are ${}_B \text{Hom}(A_B, B_B)_E$ (the right B -dual of A denoted by $(A_B)^*$) and ${}_E A_B$ with B - B -bimodule isomorphism $\text{Hom}(A_B, B_B) \otimes_E A \xrightarrow{\cong} B$ given by evaluation. Now tensor all components of eq. (21) by ${}_B (A_B)^* \otimes_E -$ and cancel $B \otimes_B$ to obtain eq. (20), where of course $P = (A_B)^* \otimes_E Q$.

($\Leftarrow$) Tensor all components of eq. (20) from the left by the natural bimodule ${}_E A_B$ given by $f \cdot a \cdot b = f(a)b$, obtain eq. (21), and reverse the argument above it. This direction of proof does not make use of generator hypothesis. $\square$

Remark 2.8. Since ${}_B A_B \oplus \Omega \cong {}_B A \otimes_B A_B$ is always the case for some B - B -bimodule Ω , the B - B -bimodules $A \otimes_B A$ and A are similar or H-equivalent under the conditions of the theorem: thus their endomorphism algebras are Morita equivalent. By Theorem 2.1, left multiplication $B \hookrightarrow E$ is depth two. There is a general Galois theory of depth two extensions which in this case specializes to total algebra $\text{End}_B A \otimes_B A_B$ and base algebra $E^B \cong \text{End}_B A_B$ as parts of a bialgebroid. It is interesting to note that base and total algebras in this case are Morita equivalent.

Let $\text{Res} = \text{Res}_H^G$ denote restriction of G -modules to H -modules in the corollary below, and $\text{Ind} = \text{Ind}_H^G$ denote induction of H -modules to G -modules.

Corollary 2.9. *A subgroup H of a finite group G is depth three if and only if*

$$\langle \text{ResIndResInd}\psi | \chi \rangle \leq n \langle \text{ResInd}\psi | \chi \rangle \quad (22)$$

for all irreducible characters ψ, χ of H .

Proof. Note that the corresponding complex group algebras $A \supseteq B$ satisfy the conditions of the theorem. One arrives at the condition on inner products of characters by tensoring a simple B -module V by the components in eq. (20). Of course, whatever simple B -module components of ${}_B A \otimes_B A \otimes_B V$ has, also ${}_B A^n \otimes_B V$ has. $\square$

For example, from the character tables of the permutation groups S_4 and S_5 [6] we compute the induction-restriction table by restricting irreducible characters on S_5 , given below in matrix form (with first column and row corresponding to trivial characters):

$$M = \begin{pmatrix} 1 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 & 0 & 1 \end{pmatrix}$$

Let $\eta_i \in \text{Irr}(H)$ and $\chi_j \in \text{Irr}(G)$ ($(i, j) \in \underline{5} \times \underline{7}$). Then from row 1, $\eta_1^G = \chi_1 + \chi_3$, $\eta_1^G \downarrow_H = 2\eta_1 + \eta_4$ and finally $\eta_1^G \downarrow_H \uparrow^G \downarrow_H = 5\eta_1 + \eta_3 + 5\eta_4 + \eta_5$. Note $\langle \eta_1^G \downarrow_H \uparrow^G \downarrow_H | \eta_3 \rangle = 1 \not\leq n \langle \eta_1^G \downarrow_H | \eta_3 \rangle = 0$ for all positive integers n , whence S_4 is not a D3 subgroup in S_5 .

Computing the 5×5 matrix $S = MM^t$, we may compute that the matrix S^3 is strictly positive, so that S_4 is a depth seven subgroup in S_5 by Theorem 2.5 and its corollary. (Observing the pattern, we might conjecture at this point that the canonical subgroup $S_n < S_{n+1}$ has depth $2n - 1$.)

2.3 DEPTH THREE QUASI-BASES

The condition (20) for a depth three extension has an interpretation in terms of split epis, including the canonical split epis of a product. This should give us depth three condition in terms of quasi-bases somewhat similar to dual bases for projective modules. Meanwhile the Frobenius hypothesis on extension $A \supseteq B$ is needed to reduce the quasi-bases to simplest terms. Suppose F is a Frobenius homomorphism $A \rightarrow B$ with dual bases $\{x_i\}$ and $\{y_i\}$ in A .

Theorem 2.10. Suppose $A \supseteq B$ is a Frobenius extension where A_B is a generator. Then $A \supseteq B$ is a depth three extension if and only if there are elements $u_i, t_i \in (A \otimes_B A \otimes_B A)^B$ such that for all $x, y \in A$,

$$x \otimes_B y = \sum_{i=1}^n t_i^1 \otimes_B t_i^2 F(t_i^3 u_i^1 F(u_i^2 F(u_i^3 x) y)) \quad (23)$$

where $u = u^1 \otimes u^2 \otimes u^3$ is Sweedler notation that suppresses a possible summation over simple tensors.

Proof. ($\Rightarrow$) First note from eq. (20) that there are mappings $f_i \in \text{Hom}({}_B A_B, {}_B A \otimes_B A_B)$ and $g_i \in \text{Hom}({}_B A \otimes_B A_B, {}_B A_B)$ such that

$$\sum_{i=1}^n f_i \circ g_i = \text{id}_{A \otimes_B A}.$$

Next recall that for any B -module M , $\text{CoInd} M \cong \text{Ind} M$ for a Frobenius extension A over B [9]; I.e., there is a natural A -module isomorphism $\text{Hom}(A_B, M_B) \cong M \otimes_B A$ via $f \mapsto \sum f(x_i) \otimes y_i$ with inverse $m \otimes a \mapsto m F(a-)$. Applied to $M = A \otimes_B A$, this restricts to $\text{Hom}({}_B A_B, {}_B A \otimes_B A_B) \cong (A \otimes_B A \otimes_B A)^B$ via $f \mapsto \sum_i f(x_i) \otimes y_i$ with inverse

$$t \mapsto t^1 \otimes t^2 F(t^3 -). \quad (24)$$

Next apply the hom-tensor relation and the Frobenius isomorphism between endomorphism ring and tensor-square of extension:

$$\begin{aligned} \text{Hom}({}_B A \otimes_B A_B, {}_B A_B) &\cong \text{Hom}(A_B, E_B)^B \\ &\cong \text{Hom}({}_B A_B, {}_B A \otimes_B A_B) \cong (A \otimes_B A \otimes_B A)^B. \end{aligned}$$

Following the isomorphisms, the forward composite mapping is given by $g \mapsto \sum_{i,j} g(x_i \otimes x_j) \otimes y_j \otimes y_i$ with inverse given by

$$u \mapsto (x \otimes y \mapsto u^1 F(u^2 F(u^3 x) y)) \quad (25)$$

for all $u \in (A \otimes_B A \otimes_B A)^B$, $x, y \in A$.

Now suppose the mappings we begin with $f_i \mapsto t_i$ and $g_i \mapsto u_i$ in $(A \otimes_B A \otimes_B A)^B$ via isomorphisms displayed above. Then eq. (23) results.

($\Leftarrow$) Define a split B - B -bimodule epimorphism $A^n \rightarrow A \otimes_B A$ by

$$(a_1, \dots, a_n) \mapsto \sum_{i=1}^n t_i^1 \otimes t_i^2 F(t_i^3 a_i) \text{ with section } A \otimes_B A \rightarrow A^n \text{ given by } x \otimes y \mapsto (u_i^1 F(u_i^2 F(u_i^3 x) y))_{i=1, \dots, n}.$$

□

For example, a left depth two quasi-bases $t_i \in (A \otimes_B A)^B$ and $\beta_i \in \text{End}_B A_B$ for $A \supseteq B$ satisfy $x \otimes y = \sum_{i=1}^n t_i \beta_i(x) y$ for all $x, y \in A$. If A is Frobenius extension of B , then $\text{End}_B A_B \cong (A \otimes_B A)^B$ via $\alpha \mapsto \sum_i \alpha(x_i) \otimes y_i$ with inverse $t \mapsto t^1 F(t^2 -)$. Let $u_i \in (A \otimes_B A)^B$ satisfy $u_i^1 F(u_i^2 -) = \beta_i$. Then

$$\left\{ \sum_j t_i^1 \otimes_B t_j^2 x_j \otimes_B y_j \right\}_{i=1, \dots, n} \quad \left\{ \sum_j x_j \otimes_B y_j u_i^1 \otimes_B u_i^2 \right\}_{i=1, \dots, n} \quad (26)$$

are D3 quasi-bases, because

$$\begin{aligned} \sum_{i,j,k} t_i^1 \otimes t_i^2 x_j F(y_j x_k F(y_k u_i^1 F(u_i^2 x) y)) &= \sum_{i,k} t_i^1 \otimes t_i^2 x_k F(y_k u_i^1 F(u_i^2 x) y) \\ &= \sum_i t_i^1 \otimes t_i^2 u_i^1 F(u_i^2 x) y = x \otimes y. \end{aligned}$$

3 HALL SUBGROUP IN FROBENIUS GROUP IS DEPTH THREE

A Frobenius group is a finite group G with nontrivial normal subgroup M (called the Frobenius kernel) which contains the centralizer of each of its nonzero elements: $C_G(\{x\}) \subseteq M$ for each $x \in M^*$ [8, 17]. This is equivalent to G having a Hall subgroup, or Frobenius complement, H such that $G = MH$, $M \cap H = \{e\}$, $H \cap H^x = \{e\}$ where $H^x = x^{-1}Hx$ for any $x \in G - H$; in addition, $M = G - \cup_{x \in G} x^{-1}H^*x$. The Hall subgroup H is not normal in G (and therefore not depth two in the terms of this paper). We will see below that $H < G$ represents a nontrivial class of examples of depth three subgroup.

For example, the permutation group S_3 is a Frobenius group with kernel $M = \langle (123) \rangle$ and $H = S_2 = \langle (12) \rangle$ or either of the two subgroups $\langle (23) \rangle$ or $\langle (13) \rangle$ are Hall subgroups.

Theorem 3.1. *Let G be a Frobenius group with Hall subgroup H . Then H is depth three subgroup of G .*

Proof. From the defining condition (22), we easily find a positive integer n if $\langle \text{ResInd}\psi | \chi \rangle > 0$ for all irreducible characters ψ, χ of H . We compute using Mackey subgroup theorem [8, p. 74] and Frobenius reciprocity, where T denotes a set of n double coset representative $\{e = g_1, g_2, \dots, g_n\}$:

$$\langle \psi^G \downarrow_H | \chi \rangle = \sum_{t \in T} \langle (\psi^t \downarrow_{H^t \cap H}) \uparrow^H, \chi \rangle = \sum_{t \in T} \langle \psi^t \downarrow_{H^t \cap H} | \chi \downarrow_{H^t \cap H} \rangle \geq n - 1$$

since $H^t \cap H = \{e\}$ for each $t \neq g_1$. Indeed it is easy to check that

$$\langle \psi^G \downarrow_H | \chi \rangle = (n - 1)(\deg \psi)(\deg \chi)$$

if $\psi \neq \chi$ and equals $1 + (n - 1)(\deg \psi)^2$ if $\chi = \psi$. □

For example the subgroup S_2 in S_3 has two double coset representatives, both irreducible characters are linear, and the values $\langle \psi^G \downarrow_H | \chi \rangle = \langle \psi^G | \chi^G \rangle$ are 1 on the off-diagonal and 2 on the diagonal, the coefficients of the matrix S in eq. (18). The proof of the theorem also follows from eq. (17), Corollary 2.3 and Mackey's theorem.

3.1 ACKNOWLEDGEMENTS

The second author is grateful to David Harbater and Gestur Olafsson for discussions related to this paper.

REFERENCES

- [1] J. L. Alperin and R.B. Bell, *Groups and Representations*, GTM **162**, Springer, New York, 1995.
- [2] S. Burciu, Coset decomposition for semisimple Hopf Algebras, *Comm. Alg.*, to appear.
- [3] S. Burciu, Normal Hopf subalgebras of a semisimple Hopf algebra, *Proc. A. M. S.*, to appear.
- [4] S. Burciu and L. Kadison, Depth two Hopf subalgebras of a semisimple Hopf algebra, *J. Algebra* **322** (2009), 162–176.

- [5] F. Castaño Iglesias and L. Kadison, Similarity, codepth two bicomodules and QF bimodules, preprint [arXiv:0712.4362](#)
- [6] W. Fulton and J. Harris, Representation Theory, GTM **129**, Springer, 1991.
- [7] F. Goodman, P. de la Harpe, and V.F.R. Jones, *Coxeter Graphs and Towers of Algebras*, M.S.R.I. Publ. **14**, Springer, Heidelberg, 1989..
- [8] I. M. Isaacs, *Character Theory of Finite Groups*, Dover, New York, 1976.
- [9] L. Kadison, *New Examples of Frobenius Extensions*, Univ. Lect. Ser. **14**, A.M.S., 1999.
- [10] L. Kadison and B. Külshammer, Depth two, normality and a trace ideal condition for Frobenius extensions, *Comm. Alg.* **34** (2006), 3103–3122.
- [11] L. Kadison, Finite depth and Jacobson-Bourbaki correspondence, *J. Pure & Applied Alg.* **212** (2008), 1822–1839.
- [12] R. G. Larson, Characters of Hopf algebras, *J. Algebra* **17** (1971), 352–368
- [13] R. G. Larson, and D. E. Radford, Finite dimensional cosemisimple Hopf Algebras in characteristic zero are semisimple, *J. Algebra* **117** (1988), 267–289.
- [14] S. Montgomery, *Hopf Algebras and Their Actions on Rings*, CBMS Regional Conf. Series in Math. Vol. 82, AMS, Providence, 1993.
- [15] W. D. Nichols and M. B. Richmond, The Grothendieck group of a Hopf algebra, I, *Comm. Alg.* **26** (1998), 1081–1095.
- [16] M. Rieffel, Category theory and quantum field theory, in: Noncommutative Rings (Berkeley, CA, 1989), 115–129, *Math. Sci. Res. Inst. Publ.* **24**, Springer, New York, 1992.
- [17] W. R. Scott, *Group Theory*, Dover, New York, 1987.
- [18] K. Szlachányi, On field algebra construction, arXiv preprint [0806.0041](#)